

## MassHealth Smoking Cessation Benefit

### **Briefing Notes**

#### Why provide a smoking cessation benefit?

Tobacco use is the leading cause of preventable death. In Massachusetts, the excess health care cost due to tobacco use is \$4.3 billion annually. <sup>1</sup>

#### In Massachusetts:

- 70 percent of adult cigarette smokers want to quit,
- 60 percent of smokers tried to quit smoking within the past year, and
- 41 percent report that they plan to quit in the next 30 days.<sup>2</sup>

Nicotine addiction from cigarette smoking is a chronic relapsing medical condition.<sup>3</sup> However, smoking cessation therapies, including counseling and medications, are available to help patients with this powerful addiction. Tobacco use treatment doubles quitting success rates.<sup>4</sup>

Coverage of tobacco use treatment increases both the use of effective treatment and the number of successful quit attempts.<sup>5</sup> Based on a study by Coffield, tobacco screening – including providing brief interventions and offering pharmacotherapy - is among the most cost-effective health insurance benefits that can be offered.<sup>6</sup>

### Timeline: Massachusetts Medicaid (MassHealth) smoking cessation benefit

June 2005	The FY 2006 Massachusetts state budget includes a requirement to provide a smoking
	cessation benefit through MassHealth to pregnant women and women with children
	under the age of three.

**April 2006** To help all MassHealth smokers quit, the state legislature mandates a MassHealth tobacco cessation benefit as part of Massachusetts Health Care Reform.

June 2006 The tobacco cessation benefit is announced to providers and members by MassHealth.

 $\label{eq:July 2006} \textbf{ Benefit is available beginning July $1^{st}$.}$ 

Aug 2006 The Massachusetts Tobacco Cessation and Prevention Program (MTCP) and MassHealth jointly distributed information about the benefit, including provider and consumer fact sheets, a consumer flyer, and website resources, to approximately 1,000 community health centers, hospitals, physician practices and state and community agencies. Local

programs funded by MTCP distribute materials in their communities.

Oct 2006 MTCP begins a radio and public transit campaign to help smokers quit. The campaign carries a tagline promoting the availability of the benefit to MassHealth members.

**Fall 2008** A University of Massachusetts survey finds a very high percentage of MassHealth subscribers are aware of the benefit.

June 2008 Originally a two-year pilot, the smoking cessation benefit becomes a permanent benefit.

2009 MTCP evaluation staff analyzes data; results show a significant increase in successful quit attempts from the MassHealth cessation benefit. The results become the basis for an article in the CDC's MMWR.



Massachusetts Department of Public Health Tobacco Cessation and Prevention Program (617) 624-5900 www.mass.gov/dph/mtcp

Make smoking history.



# MassHealth Smoking Cessation Benefit

### **Briefing Notes**

#### What the benefit covers


MassHealth members have access to tobacco cessation medications and counseling support.


- Medication coverage includes the nicotine patch, gum, lozenge, bupropion, and Chantix.
- Counseling is available one-on-one, in support groups, or by calling the Massachusetts Smokers' Helpline at 1-800-Try-To-Stop (1-800-879-8678).

To reduce barriers to utilization, neither counseling nor prior authorization is required to receive most medications; the co-pay is \$1 or \$3.

### Utilization in the first 3 years

	Pharmacotherapy				Counseling			
	FY07	FY08	FY09	FY07-09	FY07	FY08	FY09	FY07-Y09
# of Unique Users	27,692	43,341	42,253	84,215	971	1,664	2,059	4,398
(% of smokers)	(14.7%)	(23.0%)	(22.5%)	(44.8%)	(0.5%)	(0.9%)	(1.1%)	(2.3%)
# of Paid Claims	54,352	84,076	77,468	215,896	1,663	2,625	3,205	7,493
Amount of Paid Claims	\$3.945	\$7.023	\$5.906	\$16.874	\$56,233	\$106,858	\$159,841	\$288,372
	million	million	million	million				
Avg # of Claims Per User	1.94	1.91	1.81	2.47	1.70	1.55	1.55	1.69
Avg Cost Per Claim	\$73	\$84	\$76	\$78	\$34	\$41	\$50	\$38
Avg Cost Per User	\$142	\$162	\$140	\$196	\$58	\$63	\$78	\$66
# of Unique Providers	2,524	3,683	NA	NA	51	61	NA	NA





<sup>\*</sup> FFS (Fee-For-Service) / PCC (primary care clinician) plan only



Make smoking history.




## MassHealth Smoking Cessation Benefit

### **Briefing Notes**

### Results of the MassHealth smoking cessation benefit<sup>7</sup>

- Before the benefit, smoking prevalence among MassHealth adults was 38% compared to 16% in the rest of the Massachusetts adult population.
- Smoking prevalence among MassHealth adults decreased from 38% in 2006 before the benefit to 28% in 2008 (see figure).
- More than 33,000 MassHealth adults quit smoking.<sup>8</sup>
- Successful quit attempts increased significantly from 6.6% to 18.9%.



• Based on these and other findings, a short-term return on investment analysis conducted by George Washington University found that for every \$1 spent on the MassHealth smoking cessation benefit, \$3.12 was saved, resulting in a **net savings of \$2.12 for each \$1 invested**.9

Richard P, West K, Ku L (2012) The Return on Investment of a Medicaid Tobacco Cessation Program in Massachusetts. PLoS ONE. http://dx.plos.org/10.1371/journal.pone.0029665.





<sup>&</sup>lt;sup>1</sup> Huang, Xu et al. Smoking-Attributable Mortality, Morbidity, and Economic Costs (SAMMEC), Massachusetts 2006. Commonwealth of Massachusetts. March 1, 2008. Available at: <a href="http://www.mass.gov/Eeohhs2/docs/dph/tobacco">http://www.mass.gov/Eeohhs2/docs/dph/tobacco</a> control/sammec 2006.pdf. Accessed July 29, 2009.

<sup>&</sup>lt;sup>2</sup> Massachusetts Department of Public Health, Massachusetts Behavioral Risk Factor Surveillance System (BRFSS), 2007. Available at: <a href="http://www.mass.gov/Eeohhs2/docs/dph/tobacco\_control/adults\_who\_quits.pdf">http://www.mass.gov/Eeohhs2/docs/dph/tobacco\_control/adults\_who\_quits.pdf</a>. Accessed July 29, 2009.

<sup>3</sup> Jarvis MJ. ABC of smoking cessation: why people smoke. *BMJ*. 2004;328:277-279.

<sup>&</sup>lt;sup>4</sup> Fiore MC, Bailey WC, Cohen SJ, et al. *Treating Tobacco Use and Dependence: Clinical Practice Guideline*. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service; 2000.

<sup>&</sup>lt;sup>5</sup> Hopkins DP, Briss PA, Ricard CJ, et al. Task Force on Community Preventive Services. *American Journal of Preventive Medicine* 2001;20(2 Suppl):16-66.

<sup>&</sup>lt;sup>6</sup> Coffield AB, Maciosek MV, McGinnis JM, et al.. Priorities among recommended clinical preventive services. *American Journal of Preventive Medicine* 2001;21(1):1-9.

<sup>&</sup>lt;sup>7</sup> Land T, Warner D, Paskowsky M et al. (2010) Medicaid Coverage for Tobacco Dependence Treatments in Massachusetts and Associated Decreases in Smoking Prevalence. PLoS ONE 5(3): e9770. doi:10.1371/journal.pone.0009770.

<sup>&</sup>lt;sup>8</sup> A joinpoint analysis of Massachusetts BRFSS (Behavioral Risk Factor Surveillance System) data found a decrease of 10% per year from July 2006 to December 2008 (95% C.I.: 3.5% - 16.0%) and an estimated 33,000 smokers who quit.